

Cosmological phase space of R^n gravity

Alejandro Aviles Cervantes* and Jorge L. Cervantes-Cota[†]

**Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, A.P. 70-543, 04510 México D.F., México.*

[†]*Depto. de Física, Instituto Nacional de Investigaciones Nucleares, A. P. 18-1027, México D.F. 11801, México.*

Abstract. We present some exact solutions and a phase space analysis of metric $f(R)$ -gravity models of the type R^n . We divide our discussion in $n \neq 2$ and $n = 2$ models. The later model is a good approximation, at late times to the $f(R) = \frac{2}{\pi} R \tan^{-1}(R/\beta^2)$ gravity model, being this an example of a non-singular case. For $n \neq 2$ models we have found power law solutions for the scale factor that are attractors and that comply with WMAP 5-years data if $n < -2.55$ or $1.67 < n < 2$. On the other hand, the quadratic model has the de Sitter solution as an attractor, that also complies with WMAP 5-years data.

Keywords: Modified theories of gravity, $f(R)$, cosmology

PACS: 04.50.Kd, 98.80.Jk

INTRODUCTION

The standard model of cosmology (SMC), which is based upon the Friedmann-Robertson-Walker (FRW) models in the context of General Relativity (GR), is successfully to explain the main dynamical and kinematical features of our Universe, such as the stretching of light curves of type Ia supernovae at high redshifts ($z \sim 0.3 - 1$) [1, 2, 3], CMBR anisotropy measurements [4, 5], and structure formation features [6, 7], among other tests. To achieve all this it is necessary to introduce two unknown universe's components: dark matter (DM) and dark energy (DE). It is an unfortunate fact that we do not yet have any proved particle physics model that can account for these dark components, though some good candidates exist.

In the SMC one employs the Einstein-Hilbert action

$$S_{EH} = \int \frac{1}{2\kappa} R \sqrt{-g} d^4x, \quad (1)$$

where R is the Ricci scalar of the space-time, g is the determinant of the metric and $\kappa = 8\pi G$. This gives the well known Einstein field equations

$$G_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \kappa (T_{\mu\nu} + T_{\mu\nu}^{(DE)}), \quad (2)$$

where $T_{\mu\nu}$ includes all the known matter fields and, if necessary, the unknown DM. Positive acceleration is possible in virtue of the $T_{\mu\nu}^{(DE)}$ tensor, which might be a cosmological constant or a scalar field, generically called quintessence [8], to which one imposes the condition on its pressure to be negative, $p < -(1/3)\rho$. The *technology* known for the inflationary dynamics can then be applied to the recent cosmological acceleration [9].

Alternatively, in recent years it has appeared in the literature a new kind of models, generically called $f(R)$, which were initially aimed to account for the present acceleration of the universe [10, 11], hence avoiding the need of DE. It is possible that the above-mentioned observations indicate that GR needs to be modified at low energy scales that cosmologically correspond to recent times. More recently these models have been used to explain rotation curves [12, 13] and cluster dynamics without DM [14].

With the course of the years many models and properties have been published in which different fundamental variables have been chosen to vary the lagrangian, i.e., apart of the metric, one can also choose the Palatini formalism in which the connections are field variables as well, a formalism called *Palatini $f(R)$ gravity*, in contrast to the *metric $f(R)$ gravity* that only uses the metric. Both of these formalisms use matter fields independent of the connections, but one is not forced to do it so. Matter fields that possibly depend on the connections are called *metric-affine $f(R)$ gravity*; for more on this topic see [15, 16]. The present paper assumes the standard *metric $f(R)$ gravity* to analyze some exact solutions and to study the phase space. First, we study R^n gravity models with $n \neq 2$ and, secondly, we consider a non-singular $f(R)$ model that at late times is well approximated by R^2 .

$f(R)$ FIELD EQUATIONS

In order to geometrically modify GR, one can use some of the known gravity invariants, and among all possible ones, the simplest modification is to replace the lagrangian (1), $L = R$, with a generic function of the scalar curvature [17, 18], $f(R)$. Thus, the modified gravity action is generically written as

$$S = \int \frac{1}{2\kappa} f(R) \sqrt{-g} d^4x. \quad (3)$$

By performing the variation of the action one obtains the field equations of $f(R)$ theories [19]:

$$f'(R)R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}f(R) + (g_{\mu\nu}\nabla^\rho\nabla_\rho - \nabla_\mu\nabla_\nu)f'(R) = \kappa T_{\mu\nu}. \quad (4)$$

This is a coupled nonlinear partial differential equations system of fourth order in the metric. The trace of equation (4) is

$$3\nabla^\sigma\nabla_\sigma f'(R) + Rf'(R) - 2f(R) = \kappa T, \quad (5)$$

while the trace equation of GR is $R = -\kappa T$. This is a substantial difference, since equation (5) is a dynamical law of the scalar curvature, while its counterpart in GR is an algebraic law. Thus, we have introduced a new dynamical degree of freedom as we have changed the lagrangian from R to $f(R)$.

The flat cosmological Friedmann equations for $f(R)$ theories are, using $R = 6(\dot{H} + 2H^2) = 0$,

$$\frac{1}{2}f - 3(\dot{H} + H^2)f' + 18(\ddot{H}H + 4\dot{H}H^2)f'' = \kappa\rho \quad (6)$$

and

$$(\dot{H} + 3H^2)f' - \frac{1}{2}f - 2H\partial_0 f' - \partial_0 \partial_0 f' = \kappa p. \quad (7)$$

In the following we analyze these equations for some $f(R)$ models.

R^n MODELS

First, we want to analyze the simplest possible models in the $f(R)$ theories context, which are power laws of the scalar curvature,

$$f(R) = b^{-2(n-1)}R^n, \quad (8)$$

where b is a constant with units of mass and n is a nonzero real number. By using the Friedmann equation (6) we obtain

$$\frac{b^{2(n-1)}6^n}{2}(\dot{H} + 2H^2)^{n-2} \{n(n-1)\ddot{H}H + (1-n)\dot{H}^2 + (4n^2 - 7n + 4)\dot{H}H^2 + 2(2-n)H^4\} = \kappa\rho. \quad (9)$$

First we note that the case $R = 6(\dot{H} + 2H^2) = 0$ is meaningless for $n < 2$ ($n \neq 1$), in contrast to GR where $R = 0$ in vacuum or for matter fields with $T = 0$. In vacuum, with $\dot{H} + 2H^2 \neq 0$, equation (9) reduces to

$$n(n-1)\ddot{H}H + (1-n)\dot{H}^2 + (4n^2 - 7n + 4)\dot{H}H^2 + 2(2-n)H^4 = 0. \quad (10)$$

We seek for power law solutions $a(t) \propto t^\alpha$, $H = \alpha/t$. Using equation (10) we obtain the equation $2(2-n)\alpha^2 - (4n^2 - 7n + 4)\alpha + 2n^2 - 3n + 1 = 0$. One of its roots is $\alpha = 1/2$ which is a solution for any n , but it complies with $\dot{H} + 2H^2 = 0$ and for $n < 2$ ($n \neq 1$) it is not acceptable in this description. The other root is given by

$$\alpha = \frac{2n^2 - 3n + 1}{2 - n}. \quad (11)$$

For an accelerated universe, $\alpha > 1$, this equation implies $1.36 < n < 2$ or $n < -0.36$. The solution (11) is GR equivalent with a dark energy perfect fluid component with an effective equation of state parameter given by

$$w_{eff} = -1 + \frac{2(n-2)}{3(2n-1)(1-n)}. \quad (12)$$

The 5-year WMAP data constrains w_{eff} to $-0.11 < 1 + w_{eff} < 0.14$ [5]. This implies that in a power law model, with $n \neq 2$, n is constrained to $n < -2.55$ or $1.67 < n < 2$. Note as well that $w_{eff} > -1$ can only be achieved for a power law expanding universe.

By making a Legendre map of the lagrangian density from the nonlinear frame [20] -also called matter frame [11]- to the Einstein frame, Carrol et al. showed that a term




FIGURE 1. Plots v vs x . Here we show results with $v_0 > 1/2$, which asymptotically tend to power law solutions that are given by $v(x) = \text{const}$.

$1/R^m$ in the lagrangian can drive an accelerated expansion $a(t) \propto t^{(2m+1)(m+1)/(m+2)}$ [11], which is in perfect agreement with equation (11), where $n = -m$.

These power law solutions are attractors, as it can be shown by defining the variables $x = -H(t)$, $y = \dot{H}(t)$ and $v(x) = -x^2/y$ [21, 22]. Equation (10) then reads

$$n(n-1)\frac{x^5}{v^3}\frac{dv}{dx} - (2n^2 - 3n + 1)\frac{x^4}{v^2} + (4n^2 - 7n + 4)\frac{x^4}{v} + 2(n-2)x^4 = 0 \quad (13)$$

and an asymptotic behavior to a power law is identified with $v(x) \rightarrow \alpha = \text{constant}$, as $H \rightarrow 0$. By performing numerical integrations for different initial conditions v_0 we obtain the plots shown in Fig. 1 for $n = -1$ and for various values of n that are in the interval allowed by the WMAP data mentioned above. Accelerated solutions are found for $v_0 > 1/2$. Note that $v_0 = 1/2$ corresponds to $R = 0$. It can be shown that if $v_0 < 1/2$, the solutions tend to $v = 0$, i.e. $\dot{a}(t) \rightarrow 0$.

OTHER MODELS

There exist a plenty of $f(R)$ proposals [16], but have to be analyzed in order to verify they pass standard gravitational tests, such as stability criteria or to have a reliable Newtonian limit. For instance, theories with $f''(R) < 0$ and $|f''(R)| \ll 1$, such as $1/R^m$ or $\ln R$, are ruled out due to instabilities that appear when trying to model gravity fields in stars [23].

The initial $f(R)$ models, and some others, add to the lagrangian $L = R$ singular R -functions such as $-1/R^m$ (with $m > 0$) having a difficult interpretation to obtain a Newtonian limit, since as $R \rightarrow 0$ the theory diverges. We think a less restrictive type of theories is to consider $m < 0$ that are non-singular in this sense.

Motivated by the these facts we propose the following model:

$$f(R) = \frac{2}{\pi} R \tan^{-1}(R/\beta^2), \quad (14)$$

in which β is a constant with units of mass. Indeed, this model passes the stability criteria and has reliable Newtonian limit; these matters will be treated in a separate work. For $R \gg \beta^2$ the model behaves basically as $f(R) \rightarrow R - \frac{2}{\pi}\beta^2$, which is GR with a cosmological constant. We define t_β as the time when $R \sim \beta^2$ and we have a transition period. To adjust to observations we demand $\beta \sim H_0 \sim 10^{-33} \text{eV}$.

At late times, $R < \beta^2$, the model is well approximated by $f(R) \approx 2R^2/\pi\beta^2$. Using the Friedmann equation (6) in vacuum we obtain

$$2\ddot{H}H + 6\dot{H}H^2 - \dot{H}^2 = 0. \quad (15)$$

By integrating once, it yields,

$$\dot{H} + 2H^2 - CH^{1/2} = 0, \quad (16)$$

with C an integration constant. If $C = 0$ we obtain $a \propto t^{1/2}$ and $a = \text{const}$, which are not interesting for our aim. In Fig. 2 the phase space is shown for several values of C . The long-dashed curve is the equation $\dot{H} = -H^2$, and the acceleration condition is accomplished in the region where the solutions are above it. The double-dashed curve, $\dot{H} = 6H^2$, is the locus of the maximae which occur for all the solutions with $C > 0$. Accelerated solutions occur only for $C > 0$. The initial condition problem is to specify $\dot{H}_0$ and H_0 and then, by using (16), one determines $C = (\dot{H}_0 + 2H_0^2)/H_0^{1/2}$ to solve the dynamics.

In Fig. 3 we show the solution with $C = 1$. In general for $C > 0$, a phase point **A** with $H_0 > 0$ and $\dot{H}_0 < 0$ complies with $\ddot{H} > 0$, as it follows from equation (15), and the solution evolves to the phase point H_a . For a point **B** with $H_0 > 0$ and $\dot{H}_0 > 0$, $\ddot{H} > 0$ if $\dot{H} > 6H^2$ and $\ddot{H} < 0$ if $\dot{H} < 6H^2$. That is, for all $C > 0$, the solutions evolve to $H \rightarrow H_a = C^{2/3}/4^{1/3}$, that corresponds to an exponential de Sitter solution.

We can write equation (16) as $R = 6CH^{1/2}$, and we point out that if $R > 0$ the solution will have an accelerated final stage. Thus we divide the phase space in three regions:

I: $R < 0$; The solutions are never accelerated and tend to $a(t) \rightarrow a_0 = \text{cte}$.

II: The region between $R = 0$ ($\dot{H} = -2H^2$) and $\dot{H} = -H^2$. Here solutions are not accelerated but they reach t_β in a finite time and enter region III.

III: The solutions are asymptotically de Sitter, $a(t) \rightarrow a_0 e^{H_a t}$.

In order to adjust to observations we have $H_a \sim H(t_\beta) \sim H_0 \sim \beta \sim 10^{-33} \text{eV}$, and from $H(t_\beta) = C^{2/3}$ it follows that $C \sim \beta^{3/2} \sim 10^{-51} \text{eV}^{3/2}$.

Summarizing, the *Arctangent* model describes a universe which initially evolves such as the one described by GR with a cosmological constant into a stage dominated by




FIGURE 2. Phase space showing accelerated solutions for curves that are above the long-dashed curve.

a quadratic term of the scalar curvature. In this stage the solutions tend to a de Sitter space-time.

CONCLUSIONS

In the phase space analysis presented in this contribution we have analyzed R^n and the *Arctangent* model. For the former model with $n \neq 2$ we have found power law solutions that resulted to be attractors and that comply with the constraints imposed by WMAP [5] if $n < -2.55$ or $1.67 < n < 2$, yielding $w_{eff} > -1$. The *Arctangent* model, that was motivated as an example of a non-singular $f(R)$ -gravity, is approximated at late times as a quadratic function of R . The resulting equations can be once integrated, leading to the simple equation (16) with a constant (C) that depends on the initial conditions ($H_0, \dot{H}_0$). We have found three phase space regions having the following properties: In region I, that corresponds to negative values of C and hence to $R < 0$, the solutions are non-accelerating. In region II, delimited by two parabolic functions, solutions are not accelerated but they tend to region III in a finite time of the order of t_β , where solutions are accelerated and tend to a de Sitter solution. Therefore, regions II and III of the phase space end with $w_{eff} = -1$, that is also in accordance with the WMAP results.




FIGURE 3. Phase space for $C = 1$. Notice that H_a is an attractor to all solutions with $C > 0$.

ACKNOWLEDGMENTS

This work has been supported by CONACYT, grant number 84133 and PhD scholarship 215819.

REFERENCES

1. A. G. Riess et al., *A. J.*, **116**, 1009 (1998).
2. S. Perlmutter et al., *Ap. J.* **517**, 565 (1999).
3. A. Filippenko, "The Accelerating Universe and Dark Energy: Evidence from Type Ia Supernovae", in *The early Universe and observational cosmology*, edited by N. Bretón, J. L. Cervantes-Cota, M. Salgado, Springer Verlag, New York, Lect. Notes Phys., 646, 2004, pp. 191-221
4. D. N. Spergel et al., *ApJS* **170**, 377 (2007).
5. G. Hinshaw et al., *Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Data Processing, Sky Maps and Basic Results*, arXiv:0803.0732 (2008).
6. J. A. Peacock et al, 2002, in *A New Era in Cosmology* (ASP Conference Proceedings), edited by T. Shanks and N. Metcalfe, astro-ph/0204239.
7. Efstathiou G., *MNRAS* **330**, L29 (2002).
8. R.R. Caldwell, R. Dave, P. J. Steinhardt, *Phys. Rev. Lett* **80**, 1582-1585 (1998).
9. E. J. Copeland, M. Sami, S. Tsujikawa, *Int. J. Mod. Phys. D* **15**, 1753-1936 (2006).
10. S. Capozziello, *Int. J. Mod. Phys. D* **11**, 483-492 (2002).
11. S.M. Carroll, V. Duvvuri, M. Trodden, M. S. Turner, *Phys. Rev. D* **70**, 043528 (2004).
12. S. Capozziello, V. F. Cardone, S. Carloni, A. Troisi, *Phys. Lett. A* **326**, 292-296 (2004).
13. Martins C.F. and P. Salucci, *MNRAS* **381**, 1103 (2007).
14. S. Capozziello, E. De Filippis, V. Salzano, *Modelling clusters of galaxies by f(R)-gravity*, arXiv:0809.1882v1 [astro-ph] (2008).
15. T. P. Sotiriou, *Ann. Phys.* **322**, 935-966 (2007).

16. T. P. Sotiriou and V. Faraoni, *f(R) theories of gravity* (2008). arXiv:0805.1726
17. A. S. Eddington, *The Mathematical Theory of Gravity*, Cambridge U. Press, Cambridge, 1923, chapter IV.
18. G. Magnano, M. Ferraris and M. Francaviglia, *Gen. Rel. Grav.* **19**, 465 (1987).
19. H. A. Buchdahl, *MNRAS* **150**, 1-8 (1970).
20. G. Magnano and L. M. Sokolowski, *Phys. Rev. D* **50**, 5039 (1994).
21. A. A. Staborinsky, *Phys. Lett. B* **91**, 99-102 (1980).
22. S. M. Carroll et al., *Phys. Rev. D* **71**, 063513 (2005).
23. M. D. Seifert, *Rev. Mod. Phys.* **76**, 064002 (2007).